

ON SOME DESMIDS FROM KOSHI TAPPU WILDLIFE RESERVE, NEPAL

S.K. Rai and P.K. Misra*

Department of Botany, Post Graduate Campus
(Tribhuvan University) Biratnagar, Nepal

*Phycology Research Laboratory, Department of Botany
Lucknow University, Lucknow-226007, India

ABSTRACT

The present work enumerates a total 26 desmids (Chlorophyta) from ponds and reservoirs in and around Koshi Tappu Wildlife Reserve, Nepal. They belong to 8 genera i.e. *Closterium* (4), *Euastrum* (4), *Micrasterias* (3), *Cosmarium* (7), *Staurastrum* (3), *Onychonema* (2), *Hyalotheca* (1), and *Desmidium* (2). Of these, 11 taxa are new records for Nepal. Genus *Onychonema* is described here for the first time from Nepal.

Key words: Algae, chlorophyta, Desmids, Koshi Tappu, Nepal.

INTRODUCTION

The first contribution on the desmid flora of Nepal was made by Hirano (1955) who had reported 79 taxa from the Himalayan region of Western Development Region (Watanabe 1982). Later on, considerable reports have been presented by Hirano (1963, 1969, 1984), Förster (1965), Kusel-fetzmann (1969), Hickel (1973), Ichimura and Kasai (1982), Watanabe (1982, 1995), Shrestha and Manandhar (1983), Nakanishi (1986), Bando *et al.* (1989), and Habib and Chaturvedi (1995, 1997). These works seem to be sporadic and most of them were concentrated to the high altitude regions. Since then, there is no any report on the desmids published for the country within this decade. Bando *et al.* (1989) have considered that the southeastern part of Asia is the most important region for the origin and distribution of desmids. Therefore, the terai region

of Nepal, due to its hot and humid climate, favour luxuriant growth of desmids. But, literature revealed that this area has not been explored so far properly. Thus, it was felt desirable to explore the algae from Koshi Tappu area.

Koshi Tappu Wildlife Reserve (Lat. 26°34'-26°45' N, Long. 86°55'- 87°05' E; elevation 70-120 m amsl; area 175 km²) lies in the flood plain of Sapta Koshi river spreading in Sunsari and Saptari districts, more or less in rectangular shape. It was established as a protected area in 1976 and declared a Ramsar site in 1987 to conserve many endangered species including Wild buffalo (*Bubalus arnee*), Gharial (*Gavialis gangeticus*) and Gangetic dolphin (*Platanista gangetica*). The Sapta Koshi river has formed many permanent ponds and marshy lands along its length and seepage stream along the eastern embankment. The area has subtropical climate with maximum temperature in May and maximum rainfall in July.

This communication describes the morphology and taxonomy of 26 desmids belonging to 8 genera of desmidiaceae from different freshwater lentic environments of Koshi Tappu Wildlife Reserve and its adjoining area and also records their distribution in other parts in Nepal. Of these, 11 taxa viz., *Euastrum platycerum* Reinsch, *Micrasterias foliacea* Bail., *Cosmarium lundelli* Delp. var. *ellipticum* W. et G.S. West f. *minus* Prescott, *C. pardalis* Cohn., *C. subspeciosum* Nordst. var. *validius* Nordst., *C. zonatum* Lund. var. *javanicum* (Gutw.) Krieg., *Stauroastrum leptocladum* Nordst. var. *cornutum* Wille, *St. sexangulare* Lund. var. *productum* Nordst., *Onychonema filiforme* (Ehr. ex Ralfs) Roy et Biss., *O. leave* Nordst. var. *latum* W. et G.S. West, and *Desmidium baileyi* (Ralfs) Nordst. var. *baileyi* f. *tetragonum* Nordst. are new records for Nepal. *Onychonema* is a genus described here for the first time for the country. As no work exists on the desmids hitherto, all these taxa are new record for this area.

Sample No.	Date of collection	Locality
EN 96	21.12.2002	Grass green, thick layered assemblage on shallow water near Hand-pump, beside a temple in the Reserve.
EN 101	21.12.2002	Planktonic in the south-west shallow edge of Titrigachi pond.
EN 128	29.03.2003	Epiphytic on the root of <i>Eichhornia crassipes</i> and <i>Pistia stratiotes</i> on the edge of Koshi Barrage Reservoir, just north of Mahendra Highway, west from Bhanatabari.
EN 129	29.03.2003	Planktonic in the south-east shallow edge of Titrigachi pond, along the north side of embankment.

MATERIALS AND METHODS

Epiphytic forms were collected by squeezing out the root of aquatic macrophytes and planktonic forms with the help of plankton net (mesh size 50 µm). Samples were tagged, labelled and then preserved with 4% formaldehyde solution. Morphotaxonomic observation and photomicrography were done with the help of Nikon Labophot Microscope in the Phycology Research Laboratory, Department of Botany, Lucknow University, India. Identification was based on illustration and dimension of the literature mentioned below the name of each taxon.

SYSTEMATIC DESCRIPTIONS

Family: Desmidiaceae

Genus: *Closterium* Nitzsch 1817

Closterium acerosum (Schrank) Ehr. ex Ralfs (Pl. 1, Fig. 8)

Scott, A.M. and G.W. Prescott 1961, P. 9, Pl. 3, Fig. 1; Gerrath, J.F. and D.M. John 1988, P. 195, Pl. 9, Fig. 1; Prasad, B.N. and P.K. Misra 1992, P. 97, Pl. 16, Fig. 15.

Cells 145-260 µm long, 30-32 µm broad, nearly straight or slightly curved, narrowly fusiform; outer margin curved with 30-36 degrees arc, inner margin straight, gradually tapering towards the rounded-truncate or sub-acute ends, apices 4.5-6 µm broad; cell wall smooth; chloroplast 3-5 ridged with 5-6 pyrenoids in a median series in each semicell.

Sample No.: EN 96

Distribution in Nepal: Rocky wall at Pangka along Dudh Koshi river, 4600 m; shallow flow at Longponga at the end of Ngozumba glacier, 4650 m; and shallow pool at the shore of Dudh pokhari lake at Gokyu, 4750 m (Watanabe 1982); roadside ditches at Mahendranagar road and a puddle near

Mahendranagar Bazaar (Habib and Chaturvedi 1995, 1997)

Closterium diana Ehr. ex Ralfs var. *diana* (Pl. 2, Fig. 5)

Kouwets, F.A.C. 1987, P. 201, Pl. 2, Figs. 23-28; Flint, E.A. and D.B. Williamson 1998, P. 75, Pl. 2, Fig. 7

Cells 330 μm long, 26.5 μm broad; outer margin curved with 108 degree arc, inner slightly tumid or inflated in the middle, gradually tapering towards the obtusely rounded ends, apices 2-3 μm broad; cell wall smooth; chloroplast 3-4 ridged with 6-8 pyrenoids in a median series in each semicell.

Remarks: Our specimen has slightly larger dimension.

Sample No.: EN 129

Distribution in Nepal: Shallow streamlet near Begnash lake and shallow flow along rice field at Hetauda, 500 m (Watanabe 1982)

Closterium ehrenbergii Menegh. ex Ralfs (Pl. 1, Fig. 6)

Scott, A.M. and G.W. Prescott 1961, P. 11, Pl. 2, Fig. 2; Kouwets, F.A.C. 1987, P. 203, Pl. 2, Fig. 22; Gerrath, J.F. and D.M. John 1988, P. 197, Pl. 6, Figs. 1-2.

Cells 380 μm long, 59 μm broad, stout, moderately curved; outer margin with 100-125 degrees of arc, inner margin concave, slightly inflated in the middle, cells gradually tapering to the obtusely rounded ends; apices 8-10 μm broad; cell wall smooth, colourless; chloroplast 6-8 ridged with numerous scattered pyrenoids.

Sample No.: EN 101

Distribution in Nepal: Streamlet between Naubise and Mugling, 500 m, Dhading and shallow flow along rice field at Hetauda, 500 m (Watanabe 1982); rice field at Baudha, 1325 m; pool near

Pashupatinath, 1320 m; ring road near Bansbari, 1320 m, Kathmandu; pond at Dulari, 50 m, Morang; pond at Laghula, 5 km west of Itahari, 40 m; and pond at Birganj, 40 m, Parsa (Ichimura and Kasai 1982); ringroad ditches at 2 km north of Chabahil, Kathmandu (Bando *et al.* 1989); pond near Mahendranagar Bazar, Kanchanpur (Habib and Chaturvedi 1997)

Closterium setaceum Ehr. ex Ralfs (Pl. 1, Fig. 10)

Scott, A.M. and G.W. Prescott 1961, P. 13, Pl. 1, Fig. 21; Kouwets, F.A.C. 1987, P. 207, Pl. 5, Figs. 2-3; Gerrath, J.E. and D.M. John 1988, P. 204, Pl. 9, Fig. 4.

Cells 380 μm long, 13.5-15 μm broad, almost straight, median part fusiform-lanceolate with convex margins, tapering into long setaceous processes with rounded ends, apices 2-3 μm broad; cell wall longitudinally striated; chloroplast with 4-5 pyrenoids in a median series

Sample No.: EN 129

Distribution in Nepal: Pond at Dillibazar, 1300m, Kathmandu (Hirano 1963); small pond south of Rara lake, 3030 m, Mugu (Watanabe 1995)

Genus: *Euastrum* Ehrenberg 1832

Euastrum bidentatum Näg. (Pl. 2, Fig. 4)

Capdevielle, P. and A. Coute' 1980, P. 880, Pl. 2, Fig. 20; Alfinito, S. and B. Fumanti 1980, P. 880, Pl. 2, Fig. 20

Cells 44 μm long, 30 μm broad, deeply constricted, sinus slightly dilated at the extremity; isthmus narrow, 6-7 μm wide; semicells 3-lobed; polar lobe 19 μm broad with deep median incision, apical angles with a short spine; lateral lobes bilated, marginal spines not distinct; semicells with 5 protuberances, one large just above the isthmus, one on each lateral lobe and one on each side of apical notch in the polar lobe.

Sample No.: EN 101

Distribution in Nepal: Pheriche, 4200m, Khumbu as *E. bidentatum* Näg. var. *speciosum* Boldt ex Schmidle (Syn. *E. elegans* var. *speciosum* Boldt) (Förster, 1965)

Euastrum insulare (Wittr.) Roy (Pl. 1, Fig. 14)

Capdevielle, P. and A. Coute' 1980, P. 880, Pl. 2, Fig. 19; Kouwets, F.A.C. 1987, P. 216, Pl. 7, Fig. 21

Cells 21-22 μm long, 14.8 μm broad, deeply constricted, sinus narrowly linear; isthmus 4.5-5 μm wide; semicells truncate, 3 lobed; polar lobe 10 μm broad, short and broadly truncate with shallow median notch, apical angles rounded without marginal spine; lateral lobes broadly rounded; protuberances are not distinct.

Sample No.: EN 101

Distribution in Nepal: A stream at Mewa Valley (Hirano 1984)

Euastrum platycerum Reinsch (Pl. 2, Fig. 7)

Scott, A.M. and G.W. Prescott 1961, P. 33, Pl. 60, Fig. 4; Nurul Islam, A.K.M. 1970, P. 917, Pl. 17, Figs. 5-6; Gerrath, J.F. and D.M. John 1988, P. 210, Pl. 8, Figs. 13-14.

Cells 40-42 μm long, 37-37.5 μm broad, deeply constricted, sinus narrow, slightly dilated and widely open out; isthmus 8-10 μm wide; semicells 3 lobed; polar lobes 12.5-13 μm broad, truncate without median constriction, broadly rounded angles with 2 small marginal spines; lateral lobes broadly rounded with 5 small marginal spines; semicells with a rounded central protuberance just above the isthmus.

Sample No.: EN 101

Distribution in Nepal: New record for Nepal.

Euastrum spinulosum Delp. (Pl. 2, Fig. 6)

Scott, A.M. and G.W. Prescott 1961, P. 40, Pl. 10, Fig. 3; Nurul Islam, A.K.M. and A.K. Yusuf

Haroon 1980, P. 568, Pl. 22, Fig. 356; Yacubson, S. 1980, P. 301, Pl. 11, Fig. 132.

Cells 52 μm long, 47 μm broad, deeply constricted, sinus narrow, linear; isthmus 11.5-13 μm wide; semicells 5 lobed; polar lobe 17-18 μm broad, broadly truncate with shallow median notch, broadly rounded angles each with 3 small marginal spines; lateral lobes broadly rounded with 5-6 small marginal spines; cell wall granulate in the lobes; each semicell with a rounded, central protuberance.

Sample No.: EN 101

Distribution in Nepal: Ankhu Khola, 640 m and Luitel Bhanjyang, 770 m, Gorkha (Hirano 1955 as var. *inermius*); ditches at Chabahil, Kathmandu (Bando *et al.* 1989 as var. *javanicum*)

Genus: *Micrasterias* C.A. Agardh 1827

Micrasterias foliacea Bailey (Pl. 1, Fig. 2)

Turner, W.B. 1892, P. 24, Pl. 6, Figs. 12-15; Scott, A.M. and G.W. Prescott 1961, P. 48, Pl. 20, Fig. 4; Nurul Islam, A.K.M. 1970, P. 919, Pl. 9, Figs. 5-6; Pl. 12, Figs. 4-5.

Cells 60-70 μm long, 80 μm broad, united into filaments by inter-locking of polar lobes, outline rectangular, deeply constricted, sinus narrowly linear; isthmus 14-18 μm wide; semicells 5 lobed; polar lobes 23-26 μm long, 38-40 μm wide, exserted, sub-parallel base and expanded but excavated upper part, 2 unequal and asymmetrical spines at the base of excavation; lateral lobes with unequal lobelets, each lobelet terminating into truncate-emarginate apices; cell wall smooth.

Sample No.: EN 101

Distribution in Nepal: New record for Nepal.

Micrasterias mahabuleshwariensis Hobson (Pl. 2, Fig. 8)

Nurul Islam, A.K.M. 1970, P. 920, Pl. 9, Fig. 3; Prasad, B.N. and P.K. Misra 1992, P. 142, Pl. 20, Fig. 7

Cells 120 µm long, 104 µm broad, deeply constricted, sinus open with acuminate extremity; isthmus 20-21 µm wide; semicells 3-lobed in 3 symmetric planes with wide incision; polar lobes nearly subquadrate with two pairs of diverging, asymmetrical, denticulated processes; lateral lobes divided into 2 attenuated and denticulated processes by a wide, acute incision; all processes ends into 3 or 4 small spines.

Sample No.: EN 129

Distribution in Nepal: Taudaha, 1350 m, Kathmandu (Bando *et al.* 1989)

Micrasterias pinnatifida (Kütz.) Ralfs (Pl. 2, Fig. 11)

Scott, A.M. and G.W. Prescott 1961, P. 51, Pl. 12, Fig. 6; Pl. 14, Figs. 17-18; Nurul Islam, A.K.M. 1970, P. 920, Pl. 10, Figs. 3-7; Nurul Islam, A.K.M. and A.K. Yusuf Haroon 1980, P. 572, Pl. 14, Figs. 191-192, 199.

Cells 56 µm long, 60 µm broad, deeply constricted, sinus linear but open out; isthmus 10-11 µm wide; semicells 3 lobed with deep and rounded incision; polar lobe 13 µm long, 38 µm broad, nearly triangular, with shallow median notch; lateral lobes horizontal, semifusiform; all lobes acuminate toward the end with bifid spines; spines 4-5 µm long; cell wall minutely punctate.

Sample No.: EN 101

Distribution in Nepal: A pond at Luitel Bhanjyang, 770 m, Gorkha (Hirano 1955)

Genus: *Cosmarium* Corda 1834

Cosmarium javanicum Nordst. (Pl. 2, Fig. 1)

Nurul Islam, A.K.M. and A.K. Yusuf Haroon 1980, P. 576, Pl. 11, Figs. 156-157; Pl. 13, fig. 179; Bando, T. *et al.* 1989, P. 16, Fig. 7f

Cells 167 µm long, 80 µm broad, moderately constricted, sinus narrow, dilated at the apex; isthmus 40 µm wide; semicells truncate-pyramidal, basal angles rounded, sides convex, truncately rounded ends; cell wall distinctly punctate.

Sample No.: EN 101

Distribution in Nepal: A pond at Luitel Bhanjyang, 770 m, Gorkha (Hirano 1955); Taudaha, 1350 m, Kathmandu (Bando *et al.* 1989); Sharada river near Mahendranagar, Kanchanpur (Habib and Chaturvedi 1995)

Cosmarium lundelli Delp. var. *ellipticum* W. et G.S. West (Pl. 1, Fig. 11)

Scott, A.M. and G.W. Prescott 1961, P. 61, Pl. 25, Fig. 8; Prasad, B.N. and P.K. Misra 1992, P. 164, Pl. 22, Fig. 23; Nurul Islam, A.K.M. and H.M. Irfanullah 1999, P. 93, Pl. 1, Figs. 6-7.

Cells 57 µm long, 42.5 µm broad, deeply constricted, sinus narrowly linear, closed; isthmus 16 µm wide; semicells sub-semicircular with rounded basal angle and broadly rounded ends; cell wall punctate; chloroplast axile with two pyrenoids.

Sample No.: EN 101

Distribution in Nepal: A pond at Luitel Bhanjyang, 770 m, Gorkha (Hirano 1955); pond at Patandhoka, 1300 m, Lalitpur (Hirano 1963); pond near Nepal forest, Mahendranagar, Kanchanpur (Habib and Chaturvedi 1997)

Cosmarium lundelli Delp. var. *ellipticum* W. et G.S. West f. *minus* Prescott (Pl. 2, Fig. 13)

Bharati, S.G. and G.R. Hegde 1982, P. 744, Pl. 1, Fig. 6; Prasad, B.N. and P.K. Misra 1992, P. 164, Pl. 22, Fig. 23.

Cells 47 µm long, 41.5 µm broad, deeply constricted, sinus narrowly linear, slightly open out; isthmus 19.5 µm wide; semicells sub-semicircular to sub-pyramide with broadly rounded and shallow median constricted ends; cell wall punctate; chloroplast axile with two pyrenoids.

Sample No.: EN 101

Distribution in Nepal: New record for Nepal.

Cosmarium obsoletum (Hantz.) Reinsch (Pl. 2, Fig. 3)

Scott, A.M. and G.W. Prescott 1961, P. 63, Pl. 26, Fig. 1; Bharati, S.G. and G.R. Hegde 1982, P. 746, Pl. 1, Fig. 5; Kouwets, F.A.C. 1987, P. 226, Pl. 11, Fig. 15

Cells 43 µm long, 50 µm broad, transversely elliptic, deeply constricted, sinus narrowly linear with dilated apex and slightly open out; isthmus 22.5 µm wide; semicells depressed-semicircular, basal angles submamillate and slightly thickened, slightly flattened ends; cell wall punctate; chloroplast axile with two pyrenoids.

Sample No.: EN 101

Distribution in Nepal: A pond at Luitel Bhanjyang, 770 m, Gorkha (Hirano 1955)

Cosmarium pardalis Cohn. (Pl. 2, Fig. 2)

Scott, A.M. and G.W. Prescott 1961, P. 64, Pl. 29, Figs. 1-2; Nurul Islam, A.K.M. 1970, P. 926, Pl. 13, Fig. 7.

Cells 85-90 µm long, 74 µm broad, deeply constricted, sinus gradually opening from a dilated extremity; isthmus 35 µm wide; semicells reniform; cell wall granulate, granules arranged in horizontal and indistinct vertical series; chloroplast axile with 2 pyrenoids.

Sample No.: EN 128

Distribution in Nepal: New record for Nepal.

Cosmarium subspeciosum Nordst. var. *validius* Nordst. (Pl. 1, Fig. 12)

Yacubson, S. 1980, P. 300, Pl. 10, Fig. 114; Bharati, S.G. and G.R. Hegde 1982, P. 752, Pl. 9, Fig. 1; Sahin, B. 2005, P. 409, Fig. 14.

Cells 45 µm long, 32.5 µm broad, deeply constricted, sinus narrowly linear and close; isthmus 11-12 µm wide; semicells sub-rectangular or sub-pyramide with rounded angles, convex sides and very gradually attenuated towards broadly truncate ends, margins with 4 apical and 7 lateral crenations; cell wall granulate; chloroplast axile with 2 pyrenoids.

Sample No.: EN 101

Distribution in Nepal: New record for Nepal.

Cosmarium zonatum Lund. var. *javanicum* (Gutw.) Krieg. (Pl. 1, Fig. 13)

Scott, A.M. and G.W. Prescott 1961, P. 73, Pl. 28, Fig. 5 (as *C. zonatum*); Nurul Islam, A.K.M. and A.K. Yusuf Haroon 1980, P. 582, Pl. 12, Figs. 162-163.

Cells 37.5 µm long, 23.5 µm broad, sub-hexagonal, constricted, sinus linear with widely open out; isthmus 7.5-8 µm wide; semicells with broadly rounded basal angles, converging and retuse upper lateral margins, ends faintly retuse with rounded angles; cell wall thick and smooth; chloroplast with one pyrenoid.

Sample No.: EN 129

Distribution in Nepal: New record for Nepal.

Genus: *Staurostrum* Meyen 1829

Staurostrum leptocladum Nordst. var. *cornutum* Wille (Pl. 2, Fig. 12)

Turner, W.B. 1892, P. 123, Pl. 14, Figs. 12, 16; Nurul Islam, A.K.M. and A.K. Yusuf Haroon 1980, P. 590, Pl. 17, Fig. 243; Therezien, Y. 1985, P. 552, Pl. 25, Fig. 3.

Cells 36 μm long (without spines), 10-15 μm (without processes) to 55-67 μm (with processes) broad, constricted, sinus open out; isthmus 6-7 μm wide; semicells slightly elongated at the base, broadening upward, dorsal margin slightly convex with a pair of divergent spines; processes long, slightly convergent, crenulate margin with 2 terminal spines; chloroplast with one pyrenoid.

Sample No.: EN 129

Distribution in Nepal: New record for Nepal.

Staurostrum sexangulare Lund. var. *productum* Nordst. (Pl. 2, Fig. 9)

Scott, A.M. and G.W. Prescott 1961, P. 107, Pl. 46, Figs. 3-4; Nurul Islam, A.K.M. and A.K. Yusuf Haroon 1980, P. 592, Pl. 19, Fig. 289.

Cells 17.5-30 μm (without processes) and 55 μm (with processes) broad; semicells with 5 large processes, each large process provided with a small process at its lower side at the base, each process terminate into 3-5 divergent spines.

Sample No.: EN 129

Distribution in Nepal: New record for Nepal.

Staurostrum sexcostatum Bréb. ex Ralfs var. *productum* (W. West) G.S. West (Pl. 2, Fig. 10)

Kouwets, F.A.C. 1987, P. 251, Pl. 19, Fig. 7; Kim, H.S. 1996, P. 536, Pl. 7, Fig. 68.

Cells 39.5 μm long, 42-42.5 μm broad (with processes), shallow constriction with an acute

notch; isthmus 15 μm wide; semicells nearly hexagonal to cup shaped, dorsal margin straight to slightly convex; processes 6, stout, each 7.5-9 μm long, 6.3 μm broad at the base, slightly convergent with 3 marginal crenulations and 3 short terminal spines; chloroplasts with 1 pyrenoid.

Sample No.: EN 129

Distribution in Nepal: A pond at Luitel Bhanjyang, 770 m, Gorkha (Hirano 1955)

Genus: *Onychonema* Wallich 1860 (New genus)

Onychonema filiforme (Ehr. ex Ralfs) Roy et Biss. (Pl. 1, Fig. 7)

Tiffany, L.H. and M.E. Britton 1952, P. 201, Pl. 56, Figs. 621-623; Flint, E.A. and D.B. Williamson 1998, P. 95, Pl. 10, Fig. 7

Cells 15.6 μm long (without spines), 15 μm broad, united into long twisted filament by the processes, deeply constricted, sinus sublinear with acute apex and wide open out; isthmus 4.5-5 μm wide; semicells broadly elliptic or sub-reniform, dorsal and lateral margins broadly rounded, ventral margin almost straight; processes long, divergent asymmetrical with slight swelling at the ends; chloroplast axial with 1 pyrenoid.

Sample No.: EN 129

Distribution in Nepal: New record for Nepal.

Onychonema laeve Nordst. var. *latum* W. et G.S. West (Pl. 1, Fig. 5)

Turner, W.B. 1892, P. 139, Pl. 17, Fig. 15; Tiffany, L.H. and M.E. Britton 1952, P. 201, Pl. 56, Fig. 624; Scott, A.M. and G.W. Prescott 1961, P. 121, Pl. 60, Fig. 13.

Cells 15-16 μm long, 19 μm (without spines) to 22.5-23 μm (with spines) broad, deeply

constricted, sinus open, united into filaments, filament embedded in gelatinous sheath; isthmus 4-4.5 μm wide; semicells transversely elliptic; apex with 2 divergent processes, lateral angles with 1 incurved spine; chloroplast axial with 1 pyrenoid.

Sample No.: EN 129

Distribution in Nepal: New record for Nepal.

Genus: *Hyalotheca* Ehrenberg 1841

Hyalotheca dissiliens (J.E. Smith) ex Bréb. in Ralfs (Pl. 1, Fig. 3)

Tiffany, L.H. and M.E. Britton 1952, P. 204, Pl. 56, Fig. 631; Kouwets, F.A.C. 1987, P. 256, Pl. 21, Figs. 4-5.

Cells 16 μm long, 20 μm broad, cylindric to discoidal, constricted at the joints, sinus very slightly concave, united into long filament by truncate apices, filaments embedded in a gelatinous sheath; chloroplast axial with 1 pyrenoid in each semicell.

Sample No.: EN 101

Distribution in Nepal: A pond at Pisang, 3100 m, Manang (Hirano 1955); Langtang valley, 3700 m, Rasuwa (Hirano 1969); a small pond south of Rara lake, 3030 m, Mugu (Watanabe 1995)

Genus: *Desmidium* C.A. Agardh 1824

Desmidium baileyi (Ralfs) Nordst. var. *baileyi* f. *tetragonum* Nordst. (Pl. 1, Fig. 9)

Scott, A.M. and G.W. Prescott 1961, P. 124, Pl. 62, Figs. 8-9; Nurul Islam, A.K.M. and H.M. Irfanullah 1999, P. 120, Pl. 1, Fig. 1.

Cells 15-16.5 μm long, 20-20.5 μm broad, rectangular with parallel lateral margins and broad apices with deep and semi-elliptic depression, median constriction reduced into undulation, united into a straight filament without gelatinous sheath; isthmus 17.5 μm wide; cell wall smooth; chloroplast axial.

Sample No.: EN 101

Distribution in Nepal: New record for Nepal.

Desmidium swartzii (Ag.) Ag. ex Ralfs (Pl. 1, Figs. 1 and 4)

Scott, A.M. and G.W. Prescott 1961, P. 125, Pl. 63, Fig. 8; Kouwets, F.A.C. 1987, P. 256, Pl. 21, Fig. 2; Prasad, B.N. and P.K. Misra 1992, P. 204, Pl. 26, Fig. 15.

Cells 14.2-16.2 μm long, 25 μm broad, narrowly rectangular, moderately constricted, sinus linear but widely open out, united into spirally twisted filaments; isthmus 19.7-20 μm wide; semicells narrowly oblong, lateral margins obliquely truncate with upper angle protruded toward the apex, apex slightly straight without depression; cell wall smooth; chloroplast axial.

Sample No.: EN 129

Distribution in Nepal: Ringroad ditches at 2 km north of Chabahil, Kathmandu (Bando *et al.* 1989); a small pond south of Rara lake, 3030 m, Mugu (Watanabe 1995); a pond at Indo-Nepal border, Mahendranagar, Kanchanpur (Habib and Chaturvedi 1997)




Plate 1. (Figs. 1&4. *Desmidium swartzii* (Ag.) Ag. ex Ralfs, Fig. 2. *Microsterias foliacea* Bailey, Fig. 3. *Hyalotheca dissiliens* (J.E. Smith) ex Bréb. in Ralfs, Fig. 5. *Onychonema laeve* Nordst. var. *latum* W. et G.S. West, Fig. 6. *Closterium ehrenbergii* Menegh. ex Ralfs, Fig. 7. *Onychonema filiforme* (Ehr. ex Ralfs) Roy et Biss., Fig. 8. *Closterium acerosum* (Schrank) Ehr. ex Ralfs, Fig. 9. *Desmidium baileyi* (Ralfs) Nordst. var. *baileyi* f. *tetragonum* Nordst., Fig. 10. *Closterium setaceum* Ehr. ex Ralfs, Fig. 11. *Cosmarium lundelli* Delp. var. *ellipticum* W. et G.S. West, Fig. 12. *Cosmarium subspeciosum* Nordst. var. *validius* Nordst., Fig. 13. *Cosmarium zonatum* Lund. var. *javanicum* (Gutw.) Krieg., Fig. 14. *Euastrum insulare* (Wittr.) Roy)

(Scales in the figs. 3, 5, 7, 11, 12, 13 = 20 μ m; figs. 4, 9 = 30 μ m; and figs. 1, 2, 6, 8, 10, 14 = 50 μ m)




Plate 2. (Fig. 1. *Cosmarium javanicum* Nordst., Fig. 2. *Cosmarium pardalis* Cohn., Fig. 3. *Cosmarium obsoletum* (Hantz.) Reinsch, Fig. 4. *Euastrum bidentatum* Näg., Fig. 5. *Closterium diana* Ehr. ex Ralfs var. *diana*, Fig. 6. *Euastrum spinulosum* Delp., Fig. 7. *Euastrum platycerum* Reinsch, Fig. 8. *Micrasterias mahabuleshwariensis* Hobson, Fig. 9. *Staurostrum sexangulare* Lund. var. *productum* Nordst., Fig. 10. *Staurostrum sexcostatum* Bréb. ex Ralfs var. *productum* (W. West) G.S. West, Fig. 11. *Micrasterias pinnatifida* (Kütz.) Ralfs, Fig. 12. *Staurostrum leptocladum* Nordst. var. *cornutum* Wille, Fig. 13. *Cosmarium lundelli* Delp. var. *ellipticum* W. et G.S. West f. *minus* Prescott)

(Scales in the figs. 3, 4, 6, 7, 9, 10, 11, 12, 13 = 20 µm; figs. 1, 2 = 30 µm; fig. 8 = 50 µm; and fig. 5 = 80 µm)

ACKNOWLEDGEMENT

We are grateful to the Head, Department of Botany, Lucknow University, India for laboratory facility. We also acknowledge to the Director, National Botanical Research Institute, Lucknow for his consent to consult the library. Thanks to the Warden and Staffs of the Koshi Tappu Wildlife Reserve for their kind cooperation.

REFERENCES

- Alfinito, S. and B. Fumanti. 1980. Contribution to the knowledge of fresh water algae from lake Hamun-i-puzak (Iran). *Nova Hedwigia* **33**(1-4):873-882.
- Bando, T., T. Nakano and M. Watanabe. 1989. The desmid flora of Kathmandu, Nepal. *Bull. Natn. Sci. Mus., Ser. B.* Tokyo, **15**:1-25.

- Bharati, S.G. and G.R. Hegde. 1982. Desmids from Karnataka state and Goa, Part III. Genus *Cosmarium* Corda. *Nova Hedwigia* **36**:733-757.
- Capdevielle, P. and A. Coute'. 1980. Quelques *Staurostrum* Meyen (Chlorophycées, Desmidiacées) rare ou nouveaux pour la France. *Nova Hedwigia* **33**:859-882.
- Flint, E.A. and D.B. Williamson. 1998. Desmids (Chlorophyta) in two ponds in Central Canterbury, New Zealand. *Algological Studies* **91**:71-100.
- Förster, K. 1965. Beitrag zur Kenntnis der Desmidiaceen-flora von Nepal. Erg. Forschunturn Nepal Himalaya. *Khumbu Himal* **1(2)**:25-58.
- Gerrath, J.F. and D.M. John. 1988. The desmids of Ghana, West Africa-I. *Nova Hedwigia* **46(1-2)**:187-230.
- Habib, I. and U.K. Chaturvedi. 1995. On some desmids from Nepal. *J. Ind. Bot. Soc.* **74**:277-282.
- Habib, I. and U.K. Chaturvedi. 1997. Contribution to the knowledge of desmids from Nepal. *Phykos* **36**:27-36.
- Hickel, B. 1973. Limnological investigations in lakes of Pokhara valley, Nepal. *Int. Rev. ges Hydrobiol* **58(5)**:659-672.
- Hirano, M. 1955. Fresh water algae. In: *Fauna and Flora of Nepal Himalaya*. (ed.) Kihara, H. Fauna and Flora Research Society, Kyoto Univ., Japan, pp. 5-42.
- Hirano, M. 1963. Fresh water algae from the Nepal Himalaya, collected by a member of the Japanese Climbing Expedition. *Contr. Biol. Lab.*, Kyoto Univ., Japan, **16**:1-23.
- Hirano, M. 1969. Fresh water algae from Langtang Himal, Nepal Himalaya. *Contr. Biol. Lab.*, Kyoto Univ., Japan, **22**:1-42.
- Hirano, M. 1984. Fresh water algae from east Nepal. Study reported of *Baika Junior College* **32**:197-215.
- Ichimura, T. and F. Kasai. 1982. New mating groups, group H and group I of *Closterium ehrenbergii* Meneghini from Kathmandu valley and terai plains of Nepal. In: *Reports on the Cryptogamic Study in Nepal*. (ed.) Otani, Y. National Science Museum, Tokyo, Japan, pp. 61-73.
- Kim, H.S. 1996. Desmid (*Staurostrum* and *Staurodesmus*) from Kyongsangnam-Do, Korea. *Nova Hedwigia* **62(3-4)**:521-541.
- Kouwets, F.A.C. 1987. Desmids from the Auvergne (France). *Hydrobiol* **146**:193-263.
- Kusel-fetzmann, E. 1969. Einige Algen aus Nepal. *Khumbu Himal* **1(6)**:37-56.
- Nakanishi, M. 1986. Limnological study in Phewa, Begnas and Rupa lakes. In: *Studies on distribution, adaptation and evolution of microorganisms in Nepal Himalayas*. (ed.) Ishida, Y. Ministry of Education, Science and Culture, Kyoto, Japan, pp. 3-13.
- Nurul Islam, A.K.M. 1970. Contributions to the knowledge of desmids of East Pakistan, Part I. *Nova Hedwigia* **20**:903-983.
- Nurul Islam, A.K.M. and A.K. Yusuf Haroon 1980. Desmids of Bangladesh. *Int. Rev. ges Hydrobiol* **65(4)**:551-604.
- Nurul Islam, A.K.M. and H.M. Irfanullah. 1999. New records of desmids for Bangladesh- III. 24 taxa. *Bangladesh J. of Plant Taxon* **6(2)**:91-104.
- Prasad, B.N. and P.K. Misra. 1992. *Fresh water algal flora of Andaman and Nicobar Islands*. Vol. 2. B. Singh and M.P. Singh Publ., Dehradun, India, 284 p.

- Sahin, B. 2005. A preliminary checklist of desmids of Turkey. *Crypt. Algal.* **26(4)**:399-415.
- Scott, A.M. and G.W. Prescott. 1961. Indonesian desmids. *Hydrobiologia* **17(1-2)**:1-132.
- Shrestha, B. and J.D. Manandhar. 1983. Contribution to the algal flora of Kathmandu valley. *J. Inst. Sci. Techn.* **6**:1-6.
- Therezien, Y. 1985. Contribution a l'Etude des Algues d'Eau Douce de la Bolivie Les Desmidiées. *Nova Hedwigia* **41**:505-576.
- Tiffany, L.H. and M.E. Britton. 1952. *The Algae of Illinois*. Hafner Publ. Co. New York, 407 p.
- Turner, W.B. 1892. The fresh water algae of east India. *Kongl. Sv. Vet. Akademiens Handlingar* **25(5)**:1-187.
- Watanabe, M. 1982. Observation on the genus *Closterium* from Nepal. In: *Reports on the Cryptogamic Study in Nepal*. (compiled) Y. Otani. National Science Museum, Tokyo, Japan, pp. 47-59.
- Watanabe, M. 1995. Algae from lake Rara and its vicinities, Nepal Himalayas. In: *Cryptogams of the Himalayas*, Vol. 3, *Nepal and Pakistan*. (eds.) Watanabe, M. and H. Hagiwara. National Science Museum, Tsukuba, Japan, pp. 1-17.
- Yacubson, S. 1980. The phytoplankton of some fresh water bodies from Zulia state (Venezuela). *Nova Hedwigia* **33**:279-339.