

Habitat segregation between sympatric Tibetan argali *Ovis ammon hodgsoni* and blue sheep *Pseudois nayaur* in the Indian Trans-Himalaya

Tsewang Namgail^{1,2}, Joseph L. Fox^{1*} and Yash Veer Bhatnagar²

¹ Department of Biology, Faculty of Science, University of Tromsø, N-9037 Tromsø, Norway

² Wildlife Institute of India, P.O. Box 18, Dehradun – 248 001, Uttaranchal, India

(Accepted 18 June 2003)

Abstract

Tibetan argali *Ovis ammon hodgsoni* and blue sheep *Pseudois nayaur* have almost completely overlapping distributions encompassing most of the Tibetan plateau and its margins. Such a sympatric distribution of related species with similar ecological requirements implies that there is some degree of resource partitioning. This may be accomplished on the basis of habitat and/or diet separation. This study evaluated such ecological separation on the basis of physical habitat partitioning by these two sympatric ungulates in Hemis High Altitude National Park, Ladakh, India, in an area where the argali established a small new population in 1978. Such separation was tested for on the basis of expected difference between the species in their proximity to cliffs, associated with species-specific anti-predator behaviour. Tibetan argali selected habitats away from cliffs while blue sheep selected habitats close to cliffs. Blue sheep also selected steep slopes whereas argali selected gentle slopes. The two species did not differ in their use of habitats in terms of elevation. They did, however, differ in their use of plant communities; blue sheep selected sub-shrub and grass-dominated communities whilst argali selected forb-dominated communities. We suggest that the two species coexist in this site as a result of the differential use of habitat associated with their species-specific anti-predator strategies.

Key words: argali, *Ovis ammon hodgsoni*, blue sheep, *Pseudois nayaur*, habitat selection, resource partitioning, niche relationship

INTRODUCTION

The pattern of resource partitioning in ungulate communities is widely studied (Jarman, 1974; Jarman & Sinclair, 1979; Gordon & Illius, 1989; Voeten & Prins, 1999; Forsyth, 2000), but the mechanisms which produce such partitioning remain elusive. The observed pattern of resource partitioning or niche differentiation in ungulate communities could be attributed to both ecological forces (i.e. current competition and/or predation) and evolutionary history (i.e. ghost of competition past) (Connell, 1980). In coevolved communities, the extant species may have competed in the past and developed different morphological and/or behavioural characteristics through evolutionary time, thus accomplishing resource partitioning. There is however an inherent problem in invoking the so-called 'ghost of competition past', as we cannot in retrospect determine the past processes and check whether they competed in the past or not (Begon, Harper & Townsend, 1996). Therefore, the pattern of

resource partitioning in ungulate communities can only be tested and interpreted in terms of current ecological forces.

Resource partitioning studies have addressed mountain ungulate–livestock interaction for the Asiatic ibex *Capra ibex sibirica* (Bhatnagar *et al.*, 2000) and blue sheep *Pseudois nayaur* (Mishra, 2001) in the relatively dry areas of the Indian Himalayan region, but other than theoretically (Mishra *et al.*, 2002) niche separation among wild species has not been addressed in this area. Study of ecological separation among musk deer *Moschus chrysogaster*, goral *Nemorhaedus goral*, serow *Capricornis sumatraensis* and sambar *Cervus unicolor* has indicated that, in the moist and productive areas of the Greater Himalaya, these dissimilar sympatric species apparently partition the abundant resources (Green, 1987). Ungulates in the drier Trans-Himalayan region, however, live in a relatively impoverished environment (Mishra, 2001), where they are expected to compete for common resources, and coexist either by geographical or resource partitioning (Schaller, 1977). In recent years, there has been a great controversy on the relative importance of competition and predation in structuring ecological communities (see reviews by Sih *et al.*, 1985; Gurevitch,

*Corresponding author: Joseph L. Fox.
E-mail: joe.fox@ib.uit.no

Morrison & Hedges, 2000). Much of the literature has focused on competition as a plausible factor influencing differential use of resources by coexisting species (MacArthur, 1972; Pianka, 1976; Fritz, de Garine-Wichatitsky & Letessier, 1996), while importance of predation in resource partitioning is little understood. Predation may lead to resource partitioning provided the species are safest from predators in different habitats (Sih *et al.*, 1985; Repasky, 1996).

The Tibetan argali *Ovis ammon hodgsoni*, henceforth 'argali', and the blue sheep *P. nayaur* are two wild ungulates of the Bovidae subfamily Caprinae, with almost completely overlapping distributions (see Shackleton, 1997) that encompass the Tibetan plateau and its peripheral areas. Although other subspecies of *O. ammon* occur in areas without blue sheep, where overlap occurs as with *O. a. hodgsoni* such sympatric distribution of related species with similar ecological requirements implies that they avoid any high degree of competition, and that there is some degree of resource partitioning. This may be accomplished on the basis of habitat and/or diet separation. Initial food habit studies of argali and blue sheep on the Tibetan plateau (Harris & Miller, 1995; Miller & Schaller, 1998) have shown a considerable overlap in diet, suggesting that they accomplish the resource partitioning primarily by habitat separation.

Various anecdotal reports on argali and blue sheep habitat utilization suggest that although both species are denizens of mountainous areas, blue sheep prefer steep terrain near cliffs whereas argali prefer open and rolling terrain away from cliffs (Fox, Nurbu & Chundawat, 1991; Schaller, 1977, 1998). The biological basis for this difference in habitat use is explained in terms of contrasting predator avoidance strategies associated with morphological differences between the two species (Geist, 1971; Main, Weckerly & Bleich, 1996). Thus, the long slender legs of argali enhance its cursorial strategy of out-running predators on the open and rolling terrain of its preferred habitat. In contrast, the relatively short and muscular legs of the blue sheep support its agility in steep and rugged terrain (i.e. cliffs) where it retreats to avoid predators, which are generally less agile in such habitat (Bleich, 1999; Geist, 1999). Thus, based on this difference, the blue sheep and argali are expected to show some differences in their use of habitat in terms of their proximity to cliffs related to the respective anti-predator strategies. This expected difference provides the basis for the present study, especially so because no studies have hitherto actually measured argali habitat use in relation to escape terrain.

Although the species argali is widely distributed in mountainous regions of central Asia, the Tibetan argali is one of the two subspecies of *Ovis ammon* that are listed as endangered on Appendix I of the Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES). The most recent estimates suggest that, although other subspecies are much more numerous, there are < 7000 Tibetan argali left in the wild (Shackleton, 1997; Schaller, 1998). Within India there are only *c.* 200 argali remaining, mostly in Ladakh and a few

in Sikkim (Fox & Johnsingh, 1997). The small population (*c.* 20 individuals) of argali within the Hemis High Altitude National Park in Ladakh presents an interesting case where three individuals established a new population in 1978 (Fox *et al.*, 1991). The fact that these argali have not greatly increased in number since their arrival concerns the park managers.

Relying on theoretically-based expectations regarding niche (in this case habitat) separation, we have initiated an investigation of habitat use differences between this argali population and the blue sheep that share its range. This has clear implications for conservation efforts in the Hemis population, and perhaps with other argali populations. If predation was important in shaping their niche relationship, we expected blue sheep to use steep habitats near the cliffs and argali to use moderate slopes away from cliffs. Current predation does appear to be an important ecological factor in the study area, as several large predators are known to prey on these wild ungulates (Chundawat, 1992; Fox & Chundawat, 1995). Thus, we hypothesized that ecological separation between argali and blue sheep exists on the basis of physical habitat (escape terrain), and tested for this on the basis of expected difference in distance to cliff, with possible additional influences related to slope angle and/or elevation.

The study was carried out in summer (July–August) as the blue sheep, mostly males, migrate to higher elevations, where the argali occur, during this season (Chundawat, 1992). Although difference in the diet was not addressed in this summer season study, easily obtainable data on plant community type were also gathered to assess the differences in vegetation use that could be related to the aforementioned habitat differentiation.

METHODS

Study area

Our study was carried out in the north-western part of Hemis High Altitude National Park (34°N, 77°E) in the Ladakh Province of Jammu & Kashmir State, India (Fig. 1). The known range of the small argali population in the Park was delineated from local knowledge and past studies (e.g. Fox *et al.*, 1991). The entire argali range currently used encompasses *c.* 10 km² in the headwaters of the Shingo and Rumbak streams, and is situated in the Zaskar Range near the confluence of the Indus and Zaskar rivers about 20 km from Leh, the principal town of Ladakh.

Elevation within the area used by argali ranges from 4000 to 5500 m. The argali range is relatively higher than the overall blue sheep range in the whole of the Rumbak catchment (Chundawat, 1992). Within the small area of argali use, topography varies from steep, rocky slopes along the highest ridgelines to gently rolling in the mid-elevation zone and steeper again at the lowest sites. Hemis National Park is situated in the rain shadow of the Great




Fig. 1. Distribution of vegetation communities in the study area, north-western Hemis National Park, Ladakh, India.

Himalaya; hence the climate is characterized by cold and arid conditions, with low plant productivity confined to the 3-month growing season in summer (June–August). Diurnal temperatures in the Park in summer range from 15 °C to 35 °C (Chundawat, 1992). Annual precipitation in the Indus Valley at Leh is *c.* 100 mm, increasing somewhat south-westward and altitudinally to 500–1000 mm in valleys at the northern base of the Himalaya (Hartmann, 1983), and is probably *c.* 500 mm in the study area.

Vegetation of Hemis National Park is characterized by dry alpine steppe. Within the study area there are no trees, and vegetation consists of small shrubs, sub-shrubs, grasses, sedges and forbs (Fig. 1; see Namgail (2001) for a complete description of the vegetation). Predators of the argali and blue sheep include snow leopard *Uncia uncia*, wolf *Canis lupus chanco*, and wild dog *Cuon alpinus*, with red fox *Vulpes vulpes montana* and golden eagle *Aquila chrysaetos* a threat only to the young lambs.

Habitat use

Data were collected by searching for argali and blue sheep from selected trails and ridges, all with good vantage points. Observations were conducted using 8 × 40 binoculars and a 15–45X spotting scope. Records were made of species type, group size, sex, age, date and time. Individuals were considered to be solitary or belong to different groups when they stood 50 m away

from another group. Observations on argali and blue sheep were taken between 06:00 and 20:00, but most of the observations were obtained from 06:00 to 12:00, when animals were active, and effects of human-related disturbance minimal. Although not distributed evenly throughout the day, these observations can reasonably be considered a representative sample for anti-predator related habitat use. All observations were made by 1 person, thus there was no inter-observer bias.

The locations of the blue sheep and argali groups were plotted on 1:50 000 topographic sheets with elevation contour intervals of 40 m. Elevation, distance to cliff, slope angle and vegetation type (plant community) at the animal locations were recorded. Elevation was determined from the topographic sheet, and slope angle calculated from the measured distance between the nearest contour lines. Distance to cliff was determined by measuring the distance between the animal location and the nearest cliff on the map. Cliffs, very steep slopes (> 45°) on an area of >30 m diameter with vertical drops of >3 m, were identified and mapped in the field. The vegetation was classified into 5 communities, based on physiognomy and dominant species, and mapped on the topographic sheet (Fig. 1). Subsequently, the mapped blue sheep and argali locations were classified as to different vegetation communities.

As there were only 20 argali present in the area (Namgail, 2001), pseudo-replication was inevitable




Fig. 2. Locations of Tibetan argali *Ovis ammon hodgsoni* and blue sheep *Pseudois nayaur* and cliffs in the upper Rumbak and Shingo catchments of Hemis National Park, Ladakh, India.

(Machlis, Dodd & Fentress, 1985), although frequent re-grouping lessened the dependency. Owing to the high fluidity of group membership for blue sheep, and movements in and out of the study area (Namgail, 2001), it is difficult to demonstrate that the same groups shared the argali range throughout the study period. Nevertheless, a group of *c.* 27 males was frequently observed in the area through the entire study period. Blue sheep nursery groups used lower and more rugged areas in the vicinity of the study area, where they may be more secure from predators, similar to that observed for bighorn sheep *Ovis canadensis* (Bleich, Bowyer & Wehausen, 1997).

Available habitat

Availability of a habitat is the quantity of that habitat accessible to the population of animals during the study period (Manly, McDonald & Thomas, 1993). The animals were assumed to have equal access to all the available habitats as they could move across the study area within a day, and sometimes did. For availability of habitat, a systematic sample of habitat characteristics was obtained from a 1:50 000 topographic sheet of the study area. For this purpose, a point grid was overlaid on the topographic sheet and the habitat variables at the location of each point were recorded. Habitat characters (distance to cliff, slope

angle and elevation) at 366 points were sampled. These variables were determined in the same way as described for the animal habitat use.

To assess the vegetation availability, a reconnaissance survey of vegetation was conducted in the area during the study period. The plant identification was carried out in the field, using Polunin & Stainton (1985) as a reference. Once the 5 plant communities were mapped on the topographic sheet (Fig. 1), a systematic sample at the same 366 points was taken to determine their availability.

Data analysis

Habitat preference was determined by relating use of a habitat to its availability. When resources are used disproportionately to their availability, use is said to be selective (Manly *et al.*, 1993). Since individual animals were not identified and were assumed to be randomly sampled, and available population proportion of habitat characteristics was assumed to be known, the habitat selection data conformed to Design 1 with sampling protocol A, according to Manly *et al.* (1993). Although availability was sampled, since it was a systematic and intensive sampling with a point grid overlaid on the small study area, we assume that the sample available proportion represents the population proportion.

Table 1. Estimated habitat use for Tibetan argali *Ovis ammon hodgsoni* and blue sheep *Pseudois nayaur* in Hemis National Park, India. π_i , proportion of available resource units in category i ; u_i , number of used resource units in category i ; o_i , proportion of used units in category i and s_i , selection of habitat category i

Variable	π_i	Tibetan argali			Blue sheep		
		u_i	o_i	s_i	u_i	o_i	s_i
Distance to cliff (m)							
Very close to cliff (< 50)	0.119	6	0.058	–	11	0.134	0
Close to cliff (51–250)	0.374	39	0.379	0	43	0.524	+
Away from cliff (251–450)	0.336	47	0.456	+	20	0.244	0
Farther away (> 450)	0.171	11	0.107	0	8	0.098	0
Total	1.000	103	1.000		82	1.000	
Slope angle (degrees)							
Flat (< 10)	0.014	1	0.010	0	0	0.000	–
Moderate (11–30)	0.437	64	0.610	+	22	0.268	–
Steep (31–50)	0.522	40	0.381	0	57	0.695	+
Very steep (> 50)	0.027	0	0.000	–	3	0.037	0
Total	1.000	105	1.000		82	1.000	
Elevation (m)							
Low (< 4300)	0.075	4	0.038	0	2	0.024	–
Middle (4301–4600)	0.242	15	0.143	–	6	0.073	–
High (4601–4900)	0.372	31	0.295	0	33	0.402	0
Very high (> 4900)	0.311	55	0.524	+	41	0.500	+
Total	1.000	105	1.000		82	1.000	
Vegetation type							
Shrub	0.383	27	0.257	–	18	0.220	–
Sub-shrub	0.186	19	0.181	0	35	0.427	+
Grass	0.077	1	0.010	–	17	0.207	+
Sedge	0.038	0	0.000	–	4	0.049	0
Forb	0.071	57	0.543	+	1	0.012	–
Barren	0.246	1	0.010	–	7	0.085	–
Total	1.000	105	1.000		82	1.000	

+, Selection; –, avoidance; 0, use: in proportion to availability based on Bonferroni-adjusted 95% confidence intervals for habitat use.

For the determination of habitat selection, the variables were classified into distinct categories. To statistically test significant departures of use from availability, the modified χ^2 : log-likelihood chi-square test (χ_L^2) was calculated (see Manly *et al.*, 1993). If the χ_L^2 was significant, the null hypothesis: all habitats are used in proportion to their availability (no selection) was rejected. Subsequently, for each habitat category, the Bonferroni-adjusted $100(1-\alpha)\%$ confidence intervals for habitat use were constructed. A habitat category was selected if the lower confidence interval for that category was greater than the corresponding population proportion. Similarly, a habitat category was avoided when the upper confidence interval for that category excluded the corresponding population proportion.

RESULTS

A total of 82 observations on blue sheep and 105 observations on argali was recorded during the 2-month study period (Fig. 2). The argali population was composed of three adult males, 11 adult females, one yearling and five lambs (Namgail, 2001). Eighty five percent of blue sheep observations were of all-male groups, with 6% mixed groups, and 9% nursery groups.

Argali selected the habitat category ‘away from cliff’, and avoided the category ‘very close to cliff’ ($\chi_L^2 = 19.53$, $P < 0.01$). Blue sheep in contrast selected the category ‘close to cliff’ ($\chi_L^2 = 9.7$, $P < 0.05$), and its use of other categories were in proportion to the respective availabilities (Table 1). Argali also selected ‘moderate’ slopes, and avoided ‘very steep’ slopes ($\chi_L^2 = 9.26$, $P < 0.05$). Blue sheep use of ‘steep’ slopes was significantly higher than expected, and they avoided ‘moderate’ slopes ($\chi_L^2 = 29.80$, $P < 0.01$).

Argali used the ‘very high’ elevation category more than expected from its availability, and avoided the middle elevation habitats ($\chi_L^2 = 21.56$, $P < 0.01$), while it used the other two categories as expected (Table 1). Blue sheep also exhibited a similar pattern of habitat use ($\chi_L^2 = 29.80$, $P < 0.01$). Regarding vegetation, argali selected forb communities (dominated by *Potentilla* spp., *Oxytropis* spp., *Astragalus* spp., *Thermopsis* sp., *Delphinium* sp., *Dracocephalum* sp. and *Saussurea* spp.), and avoided shrub (mainly *Caragana* spp. and *Artemisia* spp.), grass (mainly *Festuca* sp., *Poa* spp. and *Elymus* spp.), and sedge (*Carex* spp. and *Kobresia* spp.) ($\chi_L^2 = 189$, $P < 0.0001$) (Table 1). Blue sheep, however, avoided forb communities, and selected sub-shrub (dominated by *Aconogonum* sp. and *Stachys* sp.) and grass communities ($\chi_L^2 = 55.0$, $P < 0.001$).

DISCUSSION

Our results support the hypothesis that ecological separation between argali and blue sheep exists on the basis of physical habitat selection. Observations of the two species showed a distinct niche divergence, with argali using relatively open terrain, and blue sheep using more rugged terrain, which conforms to previous anecdotal reports (Fox *et al.*, 1991; Schaller, 1977, 1998). Ecologists have long considered that habitat segregation serves to reduce both interference and exploitation competition, and facilitates coexistence of ecologically similar species (Pianka, 1978). Thus the habitat segregation, in terms of proximity to cliff and slope steepness, by blue sheep and argali in Hemis National Park may minimize competition and facilitate their coexistence. Although the species also differed in their association with different vegetation communities, implications regarding diet are speculative.

As there is evidence of substantial predation pressure in the study area (Chundawat, 1992), prey species may actively lower risk of predation by selecting habitat that serves as effective refuge against predators (Houtman & Dill, 1998). During the present investigation, 65% of the sightings of blue sheep were within 250 m from cliffs (escape terrain). Such high affinity by blue sheep toward cliffs, which generally support less vegetation, suggests that forage is not the only constraint in their habitat use. In contrast, 64% of the argali observations were beyond 250 m from cliffs, which illustrates the importance of open terrain in the determination of habitat use by argali. This difference in the use of habitat, in terms of proximity to cliff can be attributed to the species-specific anti-predator behaviour.

Various studies (Belovsky, 1978; Sih, 1980; Festa-Bianchet, 1988; Houtman & Dill, 1998) with their implications to the 'optimal foraging theory' suggest that an animal should forage in areas where its intake rate is highest and predation risk lowest. Although the blue sheep may be safer in the cliffs (escape terrain), there is generally less forage available in them (Namgail, 2001). Therefore, they move out of such escape terrain for feeding (Wegge, 1979). Blue sheep probably need to strike a balance between food acquisition and predator avoidance, while feeding outside the escape terrain. Their high affinity for the neighbourhood of cliffs may thus minimize overlap with argali.

Although diet competition was not addressed in this summer season study, as food is not likely to be limiting during this season (Miller & Schaller, 1998), association with plant communities was assessed. Blue sheep selected grass and sub-shrub communities, which were more abundant in vicinity of the cliffs. Argali, on the other hand, showed higher affinity for forb-dominated communities, which were more abundant in the open areas. Assuming that each species feeds on the dominant plants, such a selection pattern is at least consistent with the results of some diet-based studies (Koirala & Shrestha, 1997; Miller & Schaller, 1998) that record a preponderance of forbs in argali diet, and graminoids in blue sheep diet.

As for the nutritional relationships associated with interspecific differences in body size (Bell, 1971; Gagon & Chew, 2000), one would expect blue sheep (relatively smaller) to show a forb-dominated diet and argali a graminoid-dominated diet. The converse relationship observed in this study could be dictated by the plant community use associated with anti-predator habitat selection. In any case, because forage is not limiting at this time of year, such expectation for diet differences may not be appropriate.

The study has shown that despite the sparse vegetation (Chundawat, 1992), argali and blue sheep coexist in Hemis National Park, apparently by differential anti-predator habitat selection by the two species. The differential use of habitat as well as vegetation communities suggests little overlap on these dimensions, and presumably a reduced possibility of competition between argali and blue sheep. Although the continued presence of argali in the area thus seems possible, the slow growth of this new population may simply be the result of the limited amount of appropriate habitat here for this species. Other factors, such as possible competition with livestock may be important (Namgail, 2001); a threat that should also be assessed soon because of the precarious overall status of argali in India.

Acknowledgements

Financial support for this study was provided by the Department of Biology and the Centre for Environment and Development Studies, University of Tromsø, Norway and the Earthwatch Institute, USA. We express gratitude to Richard B. Harris and an anonymous reviewer for their helpful comments. We are grateful to Mr. Abdul Rauf Zargar, Wildlife Warden, Department of Wildlife Protection, Leh, for granting permission to work in Hemis High Altitude National Park.

REFERENCES

- Begon, M., Harper, J. L. & Townsend, C. R. (1996). *Ecology: individuals, populations and communities*. Oxford: Blackwell Scientific Publications.
- Bell, R. H. V. (1971). A grazing ecosystem in the Serengeti. *Sci. Amer.* **255**: 86–93.
- Belovsky, G. E. (1978). Diet optimisation in a generalist herbivore: the moose. *Theor. Popul. Biol.* **14**: 105–134.
- Bhatnagar, Y. V., Rawat, G. S., Johnsingh, A. J. T. & Stüwe, M. (2000). Ecological separation between ibex and resident livestock in a Trans-Himalayan protected area. In *Grassland ecology and management in protected areas of Nepal*: 71–84. Richard, C., Basent, K., Sah, J. P. & Raut, Y. (Eds). Kathmandu: International Centre for Integrated Mountain Development.
- Bleich, V. C. (1999). Mountain sheep and coyotes: patterns of predator evasion in a mountain ungulate. *J. Mammal.* **80**: 283–289.
- Bleich, V. C., Bowyer, R. T. & Wehausen, J. D. (1997). Sexual segregation in mountain sheep: resources or predation. *Wildl. Monogr.* **134**: 1–50.
- Chundawat, R. S. (1992). *The ecological studies of snow leopard and its associated prey species in Hemis National Park, Ladakh, India*. PhD thesis, University of Rajasthan.

- Connell, J. H. (1980). Diversity and the co-evolution of competitors, or ghost of competition past. *Oikos* **35**: 131–138.
- Festa-Bianchet, M. (1988). Seasonal range selection in bighorn sheep: conflicts between forage quality, forage quantity, and predator avoidance. *Oecologia (Berl.)* **75**: 580–586.
- Forsyth, D. M. (2000). Habitat selection and coexistence of the Alpine chamois (*Rupicapra rupicapra*) and Himalayan tahr (*Hemitragus jemlahicus*) in the eastern Southern Alps, New Zealand. *J. Zool. (Lond.)* **252**: 215–225.
- Fox, J. L. & Johnsingh, A. J. T. (1997). Country report for India. In *Wild sheep and goats, and their relatives*: 215–231. Shackleton, D. M. (Ed.). Cambridge: IUCN.
- Fox, J. L., Nurbu, C. & Chundawat, R. S. (1991). Tibetan argali (*Ovis ammon hodgsoni*) establish a new population. *Mammalia* **55**: 448–451.
- Fox, J. L. & Chundawat, R. S. (1995). Wolves in the Transhimalayan region of India; the continued survival of a low-density population. In *Ecology and conservation of wolves in a changing world*: 95–103. Carbyn, L. N., Fritts, S. H. & Seip, D. R. (Eds). Edmonton: University of Alberta Press.
- Fritz, H., de Garine-Wichatitsky, M. & Letessier, G. (1996). Habitat use by sympatric wild and domestic herbivores in an African savanna woodland: the influence of cattle spatial behaviour. *J. Appl. Ecol.* **33**: 589–598.
- Gagon, M. & Chew, A. E. (2000). Dietary preferences in extant African Bovidae. *J. Mammal.* **81**: 490–511.
- Geist, V. (1971). *Mountain sheep: A study of behaviour and evolution*. Chicago: University of Chicago Press.
- Geist, V. (1999). Adaptive strategies in American mountain sheep: effects of climate, latitude and altitude, ice age evolution, and neonatal security. In *Mountain sheep of North America*: 190–208. Valdez, R. & Krausman, P. R. (Eds). Tucson: University of Arizona Press.
- Gordon, I. J. & Illius, A. W. (1989). Resource partitioning by ungulates on the Isle of Rhum. *Oecologia (Berl.)* **98**: 167–175.
- Green, M. J. B. (1987). Ecological separation in Himalayan ungulates. *J. Zool. (Lond.)* **1**: 693–719.
- Gurevitch, J., Morrison, J. A. & Hedges, L. V. (2000). The interaction between competition and predation: a meta-analysis of field experiments. *Am. Nat.* **155**: 435–453.
- Harris, R. B. & Miller, J. D. (1995). Overlap in summer habitats and diets of Tibetan Plateau ungulates. *Mammalia* **59**: 197–212.
- Hartmann, H. (1983). Pflanzengesellschaften entlang der Kashmirroute in Ladakh. *Jb. Ver. Schutz der Bergwelt*: 131–137.
- Houtman, R. & Dill, L. M. (1998). The influence of predation risk on diet selectivity: a theoretical analysis. *Evol. Ecol.* **12**: 251–262.
- Jarman, P. J. (1974). The social organisation of antelope in relation to their ecology. *Behaviour* **48**: 215–267.
- Jarman, P. J. & Sinclair, A. R. E. (1979). Feeding strategy and the pattern of resource partitioning in ungulates. In *Serengeti, dynamics of an ecosystem*: 130–166. Sinclair, A. R. E. & Norton-Griffiths, M. (Eds). Chicago: University of Chicago Press.
- Koirala, R. A. & Shrestha, R. (1997). *Floristic composition of summer habitats and dietary relationships between Tibetan argali (Ovis ammon hodgsoni), naur (Pseudois nayaur) and domestic goat (Capra hircus) in the Damodar Kunda region of Upper Mustang in Nepal Himalaya*. MSc thesis, Agricultural University of Norway.
- MacArthur, R. H. (1972). *Geographical Ecology: Patterns in the distribution of species*. Princeton: Princeton University Press.
- Machlis, L., Dodd, P. W. D. & Fentress, J. C. (1985). The pooling fallacy: problems arising when individuals contribute more than one observation to the data set. *Z. Tierpsychol.* **68**: 201–214.
- Main, M. B., Weckerly, F. W. & Bleich, V. C. (1996). Sexual segregation in ungulates: new directions for research. *J. Mammal.* **77**: 449–461.
- Manly, B. F. J., McDonald, L. L. & Thomas, D. L. (1993). *Resource selection by animals*. London: Chapman and Hall.
- Miller, J. D. & Schaller, G. B. (1998). Rangeland dynamics in the Chang Tang Wildlife Reserve, Tibet. In *Karakorum-Hindukush-Himalaya: Dynamics of Change*: 125–147. Stellrecht, I. (Ed.). Koln: Rudiger Koppe Verlag.
- Mishra, C. (2001). *High altitude survival: conflicts between pastoralism and wildlife in the Trans-Himalaya*. PhD thesis, Wageningen University.
- Mishra, C., Van Wieren, S. E., Heitkönig, I. M. A. & Prins, H. H. T. (2002). A theoretical analysis of competitive exclusion in a Trans-Himalayan large-herbivore assemblage. *Anim. Conserv.* **5**: 251–258.
- Namgail, T. (2001). *Habitat selection and ecological separation between sympatric Tibetan argali and blue sheep in northern India*. MPhil thesis, University of Tromsø.
- Pianka, E. R. (1976). Competition and niche theory. In *Theoretical ecology*: 167–196. May, R. M. (Ed.). Oxford: Blackwell Scientific Publications.
- Pianka, E. R. (1978). *Evolutionary ecology*. 2nd edn. New York: Harper & Row.
- Polunin, O. & Stainton, A. (1985). *Flowers of the Himalaya*. Oxford: Oxford University Press.
- Repasky, R. R. (1996). Using vigilance behaviour to test whether predation promotes habitat partitioning. *Ecology* **77**: 1880–1887.
- Schaller, G. B. (1977). *Mountain monarchs*. Chicago: University of Chicago Press.
- Schaller, G. B. (1998). *Wildlife of the Tibetan steppe*. Chicago: University of Chicago Press.
- Shackleton, D. M. (1997). *Wild sheep and goats, and their relatives*. Cambridge: IUCN.
- Sih, A. (1980). Optimal behaviour: can foragers balance two conflicting demands? *Science* **210**: 1041–1043.
- Sih, A., Crowley, P., McPeck, M., Petranka, J. & Strohmeier, K. (1985). Predation, competition, and prey communities: a review of field experiments. *Ann. Rev. Ecol. Syst.* **16**: 269–311.
- Voeten, M. M. & Prins, H. H. T. (1999). Resource partitioning between sympatric wild and domestic herbivores in the Tarangire region of Tanzania. *Oecologia (Berl.)* **120**: 287–294.
- Wegge, P. (1979). Aspects of the population ecology of blue sheep in Nepal. *J. Asian Ecol.* **1**: 10–20.